



AutPlay® Therapy and Trauma

It is typically understood that autistic children and other neurodivergent children are more susceptible to victimization, experience traumatic effects differently than other children, and may perceive daily common events as traumatic due to their varied perceptions and sensitivity to stimuli (Guest & Ohrt, 2018). Although we know that autistic children experience trauma, it is not fully understood how traumatic stress impacts autistic individuals, and there is limited research on treatment of trauma symptoms in this population. When working on trauma with an autistic child, it is important to stay flexible and to adapt to the child's individual needs and learning styles. Also, many autistic children may present with a combination of adverse life events and psychiatric comorbidities. Neurologically, autistic children may process differently than neurotypical children. Processing speed, cognitive awareness, receptive language ability, and executive functioning ability may look very differently from working with a neurotypical or non-autistic child. These factors must be taken into consideration when attempting to address trauma issues, especially when using trauma models that have been normed on the neurotypical population such as TF-CBT and EMDR.

Adverse Childhood Experiences (ACEs) and Autism

The ACE study is an ongoing research study that explores the relationship between childhood trauma experience and long-term medical health and social consequences. Research findings suggest that autistic children encounter more ACEs than non-autistic children, and that this difference is especially pronounced in lower income families. As observed in the general population, autistic children who experience an increased number of ACEs are at elevated risk for comorbid psychiatric and medical health problems. Autistic children with an elevated number of ACEs also experience a delay in diagnosis and treatment initiation.

The core symptoms of autism may themselves predispose children to stressful and traumatic situations. For example, difficulty with social navigation could lead to increased peer rejection, bullying, and social anxiety. Experiences known to be distressing for autistic children such as unexpected schedule changes, the prevention or discouragement of repetitive or preferred behaviors, and sensory challenges, could be perceived as traumatic particularly when such distress occurs on a consistent basis, adding to the potential for comorbidity. These core symptoms would make every day social situations and new or unexpected experiences highly stressful for autistic children. It is possible that consistent rumination on stressful or traumatic experiences could lead to co-occurring symptoms of depression, anxiety, and even Post Traumatic Stress Disorder (PTSD) if a significant traumatic event has taken place.

AutPlay Therapy Approach to Trauma Work with Autistic Children

AutPlay Therapy is an integrative approach of relational, developmental, and behavioral methods. When working with autistic children and other children who are neurodivergent and are affected by trauma, a three-step progression is employed. The three-step process is further outlined in the AutPlay Therapy book – *AutPlay Therapy for Children and Adolescents on the Autism Spectrum* and in the training *Autism and Trauma: AutPlay Therapy and Beyond*. <https://courses.jentaylorplaytherapy.com/courses/autplay-trauma>

A brief explanation of the AutPlay three step process is offered below:

1) Relationship and Safety – Autistic children who have experienced trauma must begin with building relationship with the therapist - feeling safe and familiar with the therapist and the clinical office and environment is essential for trauma issues to be addressed. Therapeutic relationship should be the primary focal point of beginning therapy with strategic implementation of processes that help the child feel familiar and safe with the therapist and their surroundings.

2) Regulation – Autistic children and adolescents will need to learn to regulate their system before a formal trauma protocol can begin. This is often implemented in three phases: *Co-Facilitation* (cognitive instruction) - teaching and providing information about emotions, sensations, and what are feelings and what is regulation. *Co-Regulation* - modeling, supporting, and assisting children with moving through a series of regulation interventions and understanding regulation sensation. *Self-Regulation* – an obtained state where the child can recognize their need for regulation and understand how to access and implement tools on their own to regulate their system.

3) Emotion Recognition and Expression – Autistic children may not understand or recognize emotions, and most will have a challenging time expressing emotions in an appropriate or constructive manner. Trauma work will require helping autistic children recognize, express, and manage emotions.

It is possible that any established trauma protocol could be effective for an autistic child, but steps should be taken to ensure the above processes have been established before implementing any trauma protocol with autistic children. Professionals should not attempt to use AutPlay Therapy to treat trauma issues in autistic children without having a thorough understanding of trauma, a thorough knowledge of autism, and have been adequately trained in AutPlay Therapy protocol.

Autism and Trauma Research and Resources:

Barrows, P. (2002). Becoming verbal: Autism, trauma, and playfulness. *Journal of Child Psychotherapy*. 28(1), 53-72.

Cox, R. (2016). *The life recovery method: Autism treatment from a trauma perspective*. Robert Cox Publisher.

Earl, R. K., Peterson, J. L., Wallace, A. S., Fox, E., Ma, R., Pepper, M., & Haidar, G. (2017). Trauma and autism spectrum disorder: A resource guide. Retrieved from

<https://depts.washington.edu/hcsats/PDF/TF-%20CBT/pages/1%20Therapist%20Resources/Bernier-Lab-UW-Trauma-and-ASD-Reference-Guide-2017.pdf>

Fogler, J. M. & Phelps, R. A. (2018). *Trauma, autism, and neurodevelopmental disorders: Integrating research, practice, and policy*. Cham, Switzerland: Springer

Fuld, S. (2018). Autism spectrum disorder: The impact of stressful and traumatic life events and implications for clinical practice. *Clinical Social Work Journal*. 46(3), 210-219.

Gates, G. (2019). *Trauma, stigma, and autism: Developing resilience and loosening the grip of shame*. Philadelphia, PA: Jessica Kingsley Publishers.

Guest, J. D. & Ohrt, J. H. (2018). Utilizing child centered play therapy with children diagnosed with autism spectrum disorder and enduring trauma: A case example. *International Journal of Play Therapy*, 27(3), 157-165.

Haruvi-Lamdan, N., Horesh, D., & Golan, O. (2018). PTSD and autism spectrum disorder: Co-morbidity, gaps in research, and potential shared mechanisms. *Psychological Trauma: Theory, Research, Practice, and Policy*, 10(3), 290-299.

Hoover, D. W., Romero, E. M. G. (2019). The interactive trauma scale: A web-based measure for children with autism. *Journal of Autism and Developmental Disorders*. 49(4), 1686-1692.

Hoover, D. W., & Kaufman, J. (2018). Adverse childhood experiences in children with autism spectrum disorder. *Current opinion in psychiatry*, 31(2), 128–132.
<https://doi.org/10.1097/YCO.0000000000000390>

Hoover, D. W. (2015). The effects of psychological trauma on children with autism spectrum disorders: A research review. *Review Journal of Autism and Developmental Disorders*, 2(3), 287-299.

Kerns, C. M., Newschaffer, C. J., & Berkowitz, S. J. (2015). Traumatic childhood events and autism spectrum disorder. *Journal of Autism and Developmental Disorders*, 45(11), 3475-3486.

Kerns, C. M., Newschaffer, C. J., Berkowitz, S., & Lee, B. K. (2017). Brief report: Examining the association of autism and adverse childhood experiences in the national survey of children's health: The important role of income and co-occurring mental health conditions. *Journal of autism and developmental disorders*, 47(7), 2275–2281.
<https://doi.org/10.1007/s10803-017-3111-7>

Stack, A. & Lucyshyn, J. (2019). Autism spectrum disorder and the experience of traumatic events: Review of the current literature to inform modifications to a treatment model for children with autism. *Journal of Autism & Developmental Disorders*. 49(4), 1613-1625.